

Math 1401 - Practice Final Exam

This is a practice exam only. The actual exam may differ from this practice exam.
In fact, there are **many more questions here than will be on the final exam.**
Note: items marked **in yellow** involve concepts to be discussed on Monday.

Part 1: Integration Problems

1. Answer the following questions. Be concise:

a) State the *definition* as well as the *geometric interpretation* of:

- $f(x)$ is continuous at the point $x = a$
- the derivative of a function $f(x)$, i.e. $\frac{d}{dx} f(x)$
- the definite integral of a function $f(x)$ over an interval $[a, b]$, i.e. $\int_a^b f(x) dx$


b) What is the definition of the indefinite integral $\int f(x) dx$

c) State the first fundamental theorem of calculus. What is that theorem good for, in your own words?

d) State the second fundamental theorem of calculus. What is that theorem good for, in your own words?

e) What is the difference between $\int_a^b f(x) dx$ and $\int f(x) dx$

2. Consider the function displayed below, and state whether the indicated quantities are positive, negative, or zero.




$$\lim_{x \rightarrow -1^+} f(x)$$
$$f'(0)$$
$$f''(3)$$

$$\lim_{x \rightarrow \infty} f(x)$$
$$f'(0.5)$$
$$\int_{-1}^0 f(x) dx$$

$$f(0)$$
$$f''(0.5)$$
$$\int_{-1}^1 f(x) dx$$





3. Consider the following definite integral: $\int_0^2 4 - x^2 dx$ (the integrand is depicted below).

- a) Approximate the value of that definite integral by using 4 subdivisions and **right** rectangles in the corresponding summation.



- b) Find the exact value of that definite integral by using the first fundamental theorem of calculus. Compare with the answer in (a).

4. Below are the graphs of four functions. For each graph, decide whether $\int_0^5 f(x)dx$ is positive, negative, or zero.



5. Find the integrals (definite or indefinite) in each of the following problems, using any method you like.

$$\int 3x^2 - \frac{5}{x^3} + \sqrt[3]{x} - 2\cos(x) dx$$

$$\int x^2 + \frac{1}{\sqrt{x}} + \pi^2 dx$$

$$\int (x + \frac{1}{x})^2 dx \text{ or } \int \frac{5}{1+x^2} dx$$

$$\int x^2 + \frac{1}{x^3} + \frac{1}{x} - \cos(x) + e^x + \pi^2 dx$$

$$\int 5e^x - \frac{7}{3\sqrt{1-x^2}} dx$$

$$\int \frac{2-t^3}{\sqrt{t}} dt$$

$$\int_0^1 x^2 e^{-x^3} dx \text{ or } \int_0^1 \frac{2x-4}{x^2-4x+5} dx$$

$$\int \frac{e^x}{2+e^x} dx \text{ or } \int \frac{\ln(x)}{x} dx \text{ or } \int \frac{1}{x \ln(x)} dx$$

$$\int \frac{\ln(e^{x^2})}{e^{2\ln(x)}} dx$$

$$\int_1^3 \frac{1}{t^2} + t dt$$

$$\int_1^1 \sin(x)e^{x^2} dx \text{ or } \int_1^1 \cos(x)e^{-x^3} dx$$

$$\int_0^{\pi/2} \cos(x) - 3\sin(x) dx$$

$$\int_1^2 x(x-1) dx$$

$$\int_{-3}^3 \sqrt{9-x^2} dx \text{ (trick question)}$$

$$\int_1^4 \frac{x-1}{\sqrt{x}} dx$$

$$\int_1^2 \frac{2-t^3}{t^2} dt$$

$$\int_{\pi}^{\pi} \cos(4x^2) dx \text{ (trick question)}$$

6. Define a function $Erf(x) = \int_0^x e^{-t^2} dt$. Then find $S(0)$, find $S'(0)$, and find $S''(0)$

7. Define a function $S(x) = \int_0^x \cos^2(t^2) dt$. Then find $S(0)$, $S'(x)$, and $S''(x)$.

Part II. Review of previous concepts

Evaluate the following limits (if necessary and appropriate, use l'Hopital's Rule):

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{\cos(x)} \text{ or } \lim_{x \rightarrow 0} \frac{x}{x^2 - 1} \text{ or}$$

$$\lim_{x \rightarrow 2} \frac{x-2}{x^2 + x - 6} \text{ or } \lim_{x \rightarrow 2} \frac{x^2 - 4}{x-2}$$

$$\lim_{x \rightarrow 2} \frac{x}{x-2}$$

$$\lim_{x \rightarrow 0} \frac{\sin(3x)}{5x} \text{ or } \lim_{x \rightarrow 0} \frac{\sin(3x)}{\sin(7x)} \text{ or}$$

$$\lim_{x \rightarrow \infty} \frac{x^2 - 3x^3 + 9}{2x^3 - 6} \text{ or } \lim_{x \rightarrow \infty} \frac{x^2 - 3x^3 + 9}{2x^4 - 6} \text{ or}$$

$$\lim_{x \rightarrow \infty} \frac{\cos(3x) - 1}{x^2}$$

$$\lim_{x \rightarrow -\infty} \frac{x^2 - 3x^3 + 9}{2x^2 - 6}$$

$$\lim_{x \rightarrow \infty} \frac{x^2}{e^x} \text{ or } \lim_{x \rightarrow 0} \frac{\sin(x) - x}{x^3}$$

$$\lim_{x \rightarrow 0^+} x \ln(x)$$

$$\lim_{x \rightarrow \infty} \arctan(x)$$

$$\lim_{x \rightarrow \infty} e^{-x^2}$$

$$\lim_{x \rightarrow 1} \frac{\ln(x)}{x-1}$$

$$\lim_{x \rightarrow 0} \frac{\tan(x) - x}{x^3}$$

$$\lim_{x \rightarrow 0} \frac{\cos(x^2) - 1}{5x^4}$$

$$\lim_{x \rightarrow \pi} \frac{\sin(x)}{1 - \cos(x)}$$

$$\lim_{x \rightarrow 0^+} x \ln(x)$$

$$\lim_{x \rightarrow \frac{\pi}{2}^+} \sec(x) - \tan(x)$$

Find the number k , if any, so that the following function is continuous:

$$f(x) = \begin{cases} \frac{x^2 + 3x - 10}{x - 2} & \text{if } x \neq 2 \\ k & \text{if } x = 2 \end{cases}$$

Use the **definition** of derivative to find the derivative of $f(x) = x^2 + 2x - 2$. Do the same for $f(x) = \frac{1}{1-x}$.

Find the derivatives of the following functions, using any method you like.

$$f(x) = x^2 + \ln(x) + e^x + \pi^2$$

$$f(x) = (5x^2 + 2)^3 \cos^2(x)$$

$$f(x) = \ln(x^3)$$

$$f(x) = e^{-5x^2}$$

$$f(x) = \frac{\ln(2x+1)}{x^2+3}$$

$$f(x) = \frac{\ln(\sin(x))}{e^{-x^2}}$$

$$f(x) = \ln\left(\frac{(x+1)^2(1-x)^3}{\cos^3(x)e^x}\right)$$

$$f(x) = \tan(x^3)$$

$$f(x) = \cos(\cos^{-1}(x)) - e^{\ln(x)}$$

$$f(x) = \sin^{-1}(2x) + 3 \sec(4x)$$

$$f(x) = \arctan(x^2) (1 + x^2)^3$$

$$f(x) = \frac{(x-1)^2}{(x+1)^3} (x+2)^4$$

$$g(x) = \frac{\sqrt{x^2-1}}{x^5(x-4)^4}$$

Sketch functions like $f(x) = \frac{x^2}{x^2-1}$ or $f(x) = \frac{x}{x^2-1}$, or $f(x) = e^{-x^2}$. Or, sketch the function $f(x) = \frac{\ln(x)}{x}$ or

$$f(x) = \frac{\ln(x)}{x^2} \text{ (asymptotes } x = 0 \text{ and } y = 0).$$

Find all *relative* extrema of the function $f(x) = e^{-x^2}$ and classify them, if any.

Part III: Story Problems

- Bacteria follows exponential growth. Initially there are 100 cells but after one hour the population has increased to 420. Find the number of bacteria after 3 hours. Also find when the population reaches 10,000 cells.
- A 13 meter ladder is leaning against a wall. If the top of the ladder slips down the wall at a rate of 2 m/sec, how fast will the foot be moving away from the wall when the top is 5 m above the ground.
- A square is cut out from each corner of a rectangular piece of cardboard. The remaining sides of the cardboard are folded up to form a box without lid. If the original cardboard piece was 10 cm wide and 15 cm high, find the dimensions of the largest box possible.
- A Rectangular plot of land is fenced in with two kinds of fencing on opposite ends. Type A costs \$3 per foot, Type B costs \$2 per foot. Maximize the area if you have \$6000.00 available. x = one side of rectangle with type A fencing, y = other side of rectangle with type B fencing. Setting up the basic equations for area and cost of fencing.
- You are standing on a bridge that is 30 m above a river, and you are dropping a ball into the river. The velocity of the ball is given by the equation $v(t) = 10t$. We know that the velocity is the derivative of the distance function, hence the distance function is the antiderivative of the velocity function. Use that information to determine the distance function. Then use that distance function to determine when the ball hits the water.