



### Math 2511: Calc III - Practice Exam 3

1. State the meaning or definitions of the following terms:
  - a) vector field, conservative vector field, potential function of a vector field, volume, length of a curve, work,
  - b) curl and divergence of a vector field  $F$ , gradient of a function
  - c)  $\iint_R dA$  or  $\iint_R f(x, y) dA$
  - d)  $\iint_R dS$  or  $\int_C ds$  or  $\int_C f(x, y) ds$  or  $\int_C f(x, y) dx$  or  $\int_C f(x, y) dy$
  - e)  $\int_C \vec{F} \cdot d\vec{r}$  where  $F$  is a two or three dimensional vector field
  - f)  $\int_C M(x, y) dx + N(x, y) dy$
  - g) What does it mean when a “line integral is independent of the path”?
  - h) State the Fundamental Theorem of Line Integrals
  - i) Please state Green’s Theorem. Make sure to know when it applies, and in what situation it helps
  
2. Below are four algebraic vector fields and four sketches of vector fields. Match them.



- (1)  $F(x, y) = \langle x, y \rangle$ , (2)  $F(x, y) = \langle -y, x \rangle$ , (3)  $F(x, y) = \langle x, 1 \rangle$ , (4)  $F(x, y) = \langle 1, y \rangle$

- b) Below are two vector fields. Which one is clearly not conservative, and why?



- c) Say in the vector field [C] above you integrate over a straight line from (0,-1) to (-1,0). Is the integral positive, negative, or zero?
3. Are the following statements true or false:
- If the divergence of a vector is zero, the vector field is conservative.
  - If  $F(x, y, z)$  is a conservative vector field then  $\text{curl}(F) = 0$
  - If a line integral is independent of the path, then  $\int_C F \cdot dr = 0$  for every path C
  - If a line integral is independent of the path, then  $\int_C F \cdot dr = 0$  for every closed path C
  - If a vector field is conservative in a disk then  $\int_C F \cdot dr = 0$  for every closed path C inside that disk
  - $\iint_R dA$  denotes the surface area of the region R
  - $\iiint_R dS$  denotes the volume of the region R
  - Can you apply the Fundamental Theorem of line integrals for the function  $f(x, y, z) = xy \sin(z) \cos(x^2 + y^2)$ ?
  - Can you apply the Fundamental Theorem of line integrals for the vector field  $F(x, y) = \langle 6xy^2 - 3x^2, 6x^2y + 3y^2 - 7 \rangle$ ?
  - Can you apply Green's theorem for a curve C, which is a straight line from (0,0,0) to (1,2,3)?
4. Suppose that  $F(x, y, z) = \langle x^3y^2z, x^2z, x^2y \rangle$  is some vector field.
- Find  $\text{div}(F)$
  - Find  $\text{curl}(F)$
  - Find  $\text{curl}(\text{curl}(F))$
  - Find  $\text{div}(\text{curl}(F))$
  - $\text{grad.}$ ,  $\text{div.}$ , and  $\text{curl}$  of the vector field if appropriate for  $\langle x^2, y^2, z^2 \rangle$
  - $\text{grad.}$ ,  $\text{div.}$ , and  $\text{curl}$  of the vector field if appropriate for  $\langle \cos(y) + y \cos(x), \sin(x) - x \sin(y), xyz \rangle$
  - $\text{grad.}$ ,  $\text{div.}$ , and  $\text{curl}$  of the vector field if appropriate for  $f(x, y, z) = z \ln(x^2 + y^2)$
5. Decide which of the following vector fields are conservative. If a vector is conservative, find its potential function
- $F(x, y) = \langle 2xy, x^2 \rangle$
  - $F(x, y) = \langle e^x \cos(y), e^x \sin(y) \rangle$
  - $F(x, y, z) = \langle \sin(y), -x \cos y, 1 \rangle$
  - $F(x, y, z) = \langle 2xy, x^2 + z^2, 2zy \rangle$
  - $F(x, y) = \langle 6xy^2 - 3x^2, 6x^2y + 3y^2 - 7 \rangle$
  - $F(x, y) = \langle -2y^3 \sin(2x), 3y^2(1 + \cos(2x)) \rangle$
  - $F(x, y) = \langle 4xy + z, 2x^2 + 6y, 2z \rangle$
  - $F(x, y) = \langle 4xy + z^2, 2x^2 + 6yz, 2xz \rangle$
6. Evaluate the following integrals:
- $\iint_R \cos(x^2) dA$  where R is the triangular region bounded by  $y = 0$ ,  $y = x$ , and  $x = 1$

- b)  $\iint_R dS$ , where S is the portion of the hemisphere  $f(x, y) = \sqrt{25 - x^2 - y^2}$  that lies above the circle  $x^2 + y^2 \leq 9$
- c)  $\int_C x^2 - y + 3z ds$  where C is a line segment given by  $r(t) = \langle t, 2t, 3t \rangle$ ,  $0 \leq t \leq 1$
- d)  $\int_C F \cdot dr$  where  $F(x, y) = \langle y, x^2 \rangle$  and C is the curve given by  $r(t) = \langle 4-t, 4t-t^2 \rangle$ ,  $0 \leq t \leq 3$
- e)  $\int_C y dx + x^2 dy$  where C is a parabolic arc given by  $r(t) = \langle t, 1-t^2 \rangle$ ,  $-1 \leq t \leq 1$
- f)  $\iiint_S (x+z) dS$  where S is the first-octant portion of the cylinder  $y^2 + z^2 = 9$  between  $x = 0$  and  $x = 4$
- g) Find the flux of the vector field  $F(x, y, z) = \langle x, y, z \rangle$  through the surface given by portion of the paraboloid  $z = 4 - x^2 - y^2$  that lies above the xy-plane. Note that this surface is *not* closed.

7. For the following line integrals there is a short-cut you can use to simplify your computations (but justify your shortcut by quoting the appropriate theorem)

- a)  $\int_C F \cdot dr$  where  $F(x, y, z) = \langle e^x \cos(y), -e^x \sin(y) \rangle$  and C is the curve  $r(t) = \langle 2 \cos(t), 2 \sin(t) \rangle$ ,  $0 \leq t \leq 2\pi$
- b)  $\int_C 2xyz dx + x^2 z dy + x^2 y dz$  where C is some smooth curve from (0,0,0) to (1,4,3)
- c)  $\int_C F \cdot dr$  where  $F(x, y) = \langle y^3 + 1, 3xy^2 + 1 \rangle$  and C is the upper half of the unit circle, from (1,0) to (-1,0).
- d)  $\int_C F \cdot dr$  where  $F(x, y) = \langle y^3 x, 3xy^2 \rangle$  and C is the line segment from (-1,0) to (2,3).
- e)  $\int_C y^3 dx + (x^3 + 3xy^2) dy$  where C is the path from (0,0) to (1,1) along the graph of  $y = x^3$  and from (1,1) to (0,0) along the graph of  $y = x$ .

8. Green's Theorem

- a) Use Green's theorem to find  $\int_C F \cdot dr$  where  $F(x, y) = \langle y^3, x^3 + 3xy^2 \rangle$  and C is the circle with radius 3, oriented counter-clockwise (You may need the double-angle formula for cos somewhere during your computations)
- b) Evaluate  $\iint_R dA$  where R is the ellipse  $\frac{x^2}{4} + \frac{y^2}{9} = 1$  by using a vector field  $F(x, y) = \langle -\frac{y}{2}, \frac{x}{2} \rangle$  and the boundary C of the ellipse R.

9. Evaluate the following integrals. You can use any theorem that's appropriate:

- c)  $\int_C 2xyz dx + x^2 z dy + x^2 y dz$  where C is a smooth curve from (0,0,0) to (1,4,3)
- d)  $\int_C y dx + 2x dy$  where C is the boundary of the square with vertices (0,0), (0,2), (2,0), and (2,2)
- e)  $\int_C xy^2 dx + x^2 y dy$ , where C is given by  $r(t) = \langle 4 \cos(t), 2 \sin(t) \rangle$ , t between 0 and 2 Pi.

f)  $\int_C xy dx + x^2 dy$  where  $C$  is the boundary of the region between the graphs of  $y = x^2$  and  $y = x$ .

10. Prove the following:

a) If  $F(x, y, z) = \langle M(x, y, z), N(x, y, z), P(x, y, z) \rangle$  is any vector field where  $M, N, P$  are twice continuously differentiable then  $\operatorname{div}(\operatorname{curl}(F)) = 0$

b) A function (not a vector field)  $f(x, y, z)$  is called harmonic if  $\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = 0$ . Show that for

any function  $f(x, y, z)$  the function  $\frac{1}{f(x, y, z)}$  is harmonic.

c) Use Green's Theorem to prove that integrals of a conservative vector fields over closed curves are zero (assuming that the closed curve encloses a simply connected region and all conditions of Green's theorem are satisfied).