







## Calc 3 – Final Exam

This is a take-home, open-book, open notes exam; you even may use Maple or Wolfram Alpha to assist you in calculations. You must, however, complete it *entirely on your own*. It is due on the *last* day of exams – no exception. Please indicate clearly where each problem starts and do not forget to put your name on your exam.

1. Please state the following:
  - a) an equation relating the dot product of two vectors with the angle between them
  - b) the definition of the gradient of a function  $f(x, y, z)$  and its properties
  - c) the Divergence Theorem (also known as Gauss' Theorem)
  
2. Match the following pictures with the algebraic expressions below.



- (1)  $f(x, y) = 6 - x^2 - y^2$       (2)  $f(x, y) = y^2 - x^2$       (3)  $r(t) = \langle \cos(t), \sin(t), t \rangle$   
 (4)  $r(t) = \langle \cos(t), t, \sin(t) \rangle$       (5)  $F(x, y) = \langle x, 1 \rangle$       (6)  $F(x, y) = \langle 1, y \rangle$

3. Determine if the plane through the points  $P(1,0,0)$ ,  $Q(0,2,0)$ , and  $R(0,0,3)$  is perpendicular to the plane given by the equation  $2x - 2y - 3z = 1$
  
4. A baseball is hit 4 feet above ground at an initial velocity of  $\langle 80, 80 \rangle$  feet per second. Find the maximum height reached by the baseball. Will it clear a 15-foot high fence located 350 feet from home base?
  
5. Determine the following limits, if possible, or explain why they don't exist.

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x+2y+3}{x^2+2y^2+3}$$

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2-y^2}{x^2+y^2}$$

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2+y^4}$$

6. Find all critical points and test them for relative extrema for the function  $f(x, y) = -x^3 + 3xy - \frac{3}{2}y^2$   
(Hint: There are two critical points)
7. Which of the following vector fields is conservative? If it is conservative, find the potential function:
- $F(x, y) = \langle 2xy + y^2 - y, x^2 + 2xy - x \rangle$
  - $F(x, y) = \langle y \sin(x^2), x \cos(y^2) \rangle$
  - $F(x, y, z) = \langle 2xy + yz - 2xz^2, x^2 + xz, xy - 2zx^2 \rangle$
8. Evaluate the following integrals:
- $\iint_R e^{x^2} dA$ , where  $R$  is the triangular region bounded by  $y = 0$ ,  $y = x$ , and  $x = 1$
  - $\int_C x^2 + y^2 ds$ , where  $C$  is the curve given by  $r(t) = \langle 2 \cos(t), 2 \sin(t) \rangle$  for  $0 \leq t \leq \pi$
  - $\int_C \vec{F} d\vec{r}$ , where  $\vec{F}(x, y) = \langle -y, x \rangle$  and  $C$  is the line segment from  $P(-1, -1)$  to  $Q(2, 3)$
  - The flux of the vector field  $\vec{F}(x, y, z) = \langle x, y, z \rangle$ , where  $S$  is the portion of the surface  $z = 10 - 2x - 2y$  between the coordinate planes.
9. For the following integrals there are at least two ways to evaluate them. Use the most convenient method and quote the appropriate theorem.
- $\int_C \vec{F} \cdot d\vec{r}$  where  $\vec{F}(x, y) = \langle 2xy^3 + x \cos(x), 3x^2y^2 - \sin(y)e^y \rangle$  and  $C$  is the closed curve given by the boundary of the square with corner points  $(-1, -1)$ ,  $(-1, 1)$ ,  $(1, -1)$ , and  $(1, 1)$ .
  - $\int_C \vec{F} d\vec{r}$  where  $F(x, y) = \langle xy^2, x^2y \rangle$  where  $C$  is a curve from  $(-1, 0)$  to  $(1, 2)$
  - $\int_C (3x^2y - y^2)dx + x^3dy$  where  $C$  is the closed curve given by the boundary of the triangle with corner points  $(0, 0)$ ,  $(0, 1)$ , and  $(1, 0)$ , oriented counter-clockwise.
  - $\iint_S \vec{F} \cdot \vec{n} dS$  where  $F(x, y, z) = \langle x, 2y, 3z \rangle$  and  $S$  is given by  $x^2 + y^2 + z^2 = 9$
  - $\int_C \vec{F} dr$  where  $F(x, y, z) = \langle z^2, x^2, y^2 \rangle$  and  $C$  is the boundary of the surface  $S$  given by  $z = 1 - x - y$  between the coordinate planes, oriented counter-clockwise.