

### Math 2511: Calc III - Practice Exam 3

1. State the meaning or definitions of the following terms:

- a) vector field, conservative vector field, potential function of a vector field, volume, length of a curve, work, surface area
- b) curl and divergence of a vector field  $F$ , gradient of a function
- c)  $\iint_R dA$  or  $\iint_R f(x, y)dA$  or  $\iiint_Q f(x, y, z)dV$
- d)  $\int_C ds$  or  $\int_C f(x, y)ds$  or  $\int_C f(x, y)dx$  or  $\int_C f(x, y)dy$
- e)  $\int_C \vec{F} \cdot d\vec{r}$
- f)  $\iint_S g(x, y, z) \cdot dS$
- g)  $\int_C M(x, y, z)dx + N(x, y, z)dy + P(x, y, z)dz$
- h) What does it mean when a "line integral is independent of the path"?

see notes or book





i) State the Fundamental Theorem of Line Integrals. Make sure to know when it applies, and when it helps.]

$\int_C \vec{F} \cdot d\vec{r} = f(B) - f(A)$ ,  $C$  curve from  $A$  to  $B$ ,  $f$  potential of  $\vec{F}$ . Applies if  $\vec{F}$  is conservative

j) State Green's Theorem. Make sure to know when it applies, and in what situation it helps.



$\oint_C Mdx + Ndy = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dA$ ,  $C$  closed curve,  $R$  inside of  $C$  pos. oriented

2. Below are four algebraic vector fields and four sketches of vector fields. Match them.



- (1)  $F(x, y) = \langle x, y \rangle$ , (2)  $F(x, y) = \langle -y, x \rangle$ , (3)  $F(x, y) = \langle x, 1 \rangle$ , (4)  $F(x, y) = \langle 1, y \rangle$

b) Below are two vector fields. Which one is clearly not conservative, and why?



The first because  $\oint_C \vec{F} \cdot d\vec{r} \neq 0$

c) Say in the left vector field above you integrate over a straight line from  $(0, -1)$  to  $(1, 0)$ . Is the integral positive, negative, or zero?

pos (out work)

How about if you integrate from  $(-2,1)$  to  $(2,1)$ ? *neg. (cosh work)*

How about from  $(-2,-1)$  to  $(2,-1)$ ? *positive*

3. Are the following statements true or false:

a) If the divergence of a vector is zero, the vector field is conservative.  $\nabla$

b) If  $F(x, y, z)$  is a conservative vector field then  $\text{curl}(F) = 0$   $\nabla$

c) If a line integral is independent of the path, then  $\int_C F \cdot dr = 0$  for every path  $C$   $\nabla$

d) If a vector field is conservative then  $\int_C F \cdot dr = 0$  for every closed path  $C$   $\nabla$

e)  $\iint_R dA$  denotes the surface area of the region  $R$   $\nabla$

f)  $\iint_R f(x, y) dA$  denotes the volume of the region under the surface  $f(x, y)$  and over  $R$ , if  $f$  is positive.  $\nabla$

g) Can you apply the Fundamental Theorem of line integrals for the function  $f(x, y, z) = xy \sin(z) \cos(x^2 + y^2)$ ?  $\nabla$   
*applies to vector fields*

h) Can you apply the Fundamental Theorem of line integrals for the vector field  $\nabla$   
 $F(x, y) = \langle 6xy^2 - 3x^2, 6x^2y + 3y^2 - 7 \rangle$ ? *is conservative*

i) Can you apply Green's theorem for a curve  $C$ , which is a straight line from  $(0,0,0)$  to  $(1,2,3)$ ?  $\nabla$   
*requires closed curve*

4. Suppose that  $F(x, y, z) = \langle x^3y^2z, x^2z, x^2y \rangle$  is some vector field.

a) Find  $\text{div}(F)$

$$F_x + F_y + F_z = 3x^2y^2z + 0 + 0$$

b) Find  $\text{curl}(F)$

$$\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^3y^2z & x^2z & x^2y \end{vmatrix} = \langle x^2 - x^2, (2xy - x^2y^2), 2xz - 2x^2yz \rangle$$

c) Find  $\text{curl}(\text{curl}(F))$

*do curl again ...*

d) Find  $\text{div}(\text{curl}(F))$

$$\text{div}(0, x^3y^2 - 2xy, 2xz - 2x^2yz) = 0 + 2x^2y + 2x - 2x^2y = \underline{2x}$$

e)  $\text{grad.}$ ,  $\text{div.}$ , and  $\text{curl}$  of the vector field if appropriate for  $\langle x^2, y^2, z^2 \rangle$

$$\text{div}(x^2\hat{i} + y^2\hat{j} + z^2\hat{k}) = 2(x+y+z), \quad \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^2 & y^2 & z^2 \end{vmatrix} = \langle 0, 0, 0 \rangle$$

f)  $\text{grad.}$ ,  $\text{div.}$ , and  $\text{curl}$  of the vector field if appropriate for  $\langle \cos(y) + y\cos(x), \sin(x) - x\sin(y), xyz \rangle$

*div and curl are appropriate, left as HW*

g) grad., div., and curl of the vector field if appropriate for  $f(x, y, z) = z \ln(x^2 + y^2)$

$$\text{grad} (z \ln(x^2 + y^2)) = \left\langle \frac{2xz}{x^2 + y^2}, \frac{2yz}{x^2 + y^2}, \ln(x^2 + y^2) \right\rangle$$

5. Decide which of the following vector fields are conservative. If a vector is conservative, find its potential function

a)  $F(x, y) = \langle 2xy, x^2 \rangle$

conservative,  $f = x^2 y + C$

b)  $F(x, y) = \langle e^x \cos(y), e^x \sin(y) \rangle$

not conservative

c)  $F(x, y, z) = \langle \sin(y), -x \cos y, 1 \rangle$

d)  $F(x, y, z) = \langle 2xy, x^2 + z^2, 2zy \rangle$

e)  $F(x, y) = \langle 6xy^2 - 3x^2, 6x^2 y + 3y^2 - 7 \rangle$

conservative;  $f = 2x^2 y^2 + x^3 + y^3 - 7y + C$

f)  $F(x, y) = \langle -2y^3 \sin(2x), 3y^2(1 + \cos(2x)) \rangle$


not

g)  $F(x, y, z) = \langle 4xy + z, 2x^2 + 6y, 2z \rangle$

h)  $F(x, y, z) = \langle 4xy + z^2, 2x^2 + 6yz, 2xz \rangle$

6. Evaluate the following integrals:

a)  $\iint_R \cos(x^2) dA$  where R is the triangular region bounded by  $y = 0$ ,  $y = x$ , and  $x = 1$



$$\int_0^1 \int_0^x \cos(x^2) dy dx = \int_0^1 x \cos(x^2) dx = \frac{1}{2} \sin(x^2) \Big|_0^1 = \frac{1}{2} \sin(1)$$

b)  $\int_0^1 \int_1^{2y} x^2 y^3 dx dy$  use computer

$$r' = \langle 2t, 1 \rangle$$

c)  $\int_C ds$ , where C is the curve given by  $r(t) = \langle t^2, 1+t \rangle$ ,  $0 \leq t \leq 2$  (you might want to use Maple at some point)

$$\int_0^2 \sqrt{4t^2 + 1} dt =$$

use computer

d)  $\int_C x^2 y^3 dx$ , where C is the curve given by  $r(t) = \langle t^2, t^3 \rangle$ ,  $0 \leq t \leq 2$

$$\int_0^2 (t^4)^2 (t^3)^3 2t dt =$$

use computer

e)  $\int_C x^2 - y + 3z ds$  where C is the circle  $r(t) = \langle 2\cos(t), 2\sin(t) \rangle$ ,  $0 \leq t \leq \pi$

$$\int_0^\pi (2\cos(t))^2 - (2\sin(t)) + 3\sqrt{4(\cos^2 + \sin^2)} dt =$$

computer

f)  $\int_C x^2 - y + 3z ds$  where C is a line segment given by  $r(t) = \langle t, 2t, 3t \rangle$ ,  $0 \leq t \leq 1$

$$\int_0^1 (t)^2 - (2t) + 3(3t) \cdot \sqrt{1+4+9} dt =$$

computer

g)  $\int_C F \cdot dr$  where  $F(x, y) = \langle y, x^2 \rangle$  and C is the curve given by  $r(t) = \langle 4-t, 4t-t^2 \rangle$ ,  $0 \leq t \leq 3$

$$\int_C y dx + x^2 dy = \int_0^3 (4-t-t^2)(-1) dt + (4-t)^2 (4-2t) dt =$$

computer

h)  $\int_C F \cdot dr$  where  $F(x, y, z) = \langle yz, x^2, zy \rangle$  and C is the curve given by  $r(t) = \langle 1-t, 3t, 2-t^2 \rangle$ ,  $1 \leq t \leq 3$

$$\int_C yz dx + x^2 dy + zy dz = \int_1^3 (3t)(2-t^2)(-1) dt + (1-t)^2 3 dt + (2-t^2)(3t)(-2t) dt =$$

computer

i)  $\int_C y dx + x^2 dy$  where C is a parabolic arc given by  $r(t) = \langle t, 1-t^2 \rangle$ ,  $-1 \leq t \leq 1$

$$\int_{-1}^1 (1-t^2) dx + (t)^2 (-2t) dt =$$

computer

- j) Find the surface integral  $\iint_S x - 2y + z dS$ , where S is the surface  $z = 10 - 2x + 2y$  such that x is between 0 and 2 and y is between 0 and 4.

$$\int_0^2 \int_0^4 x - 2y + (10 - 2x + 2y) \sqrt{1 + 4 + 4} dy dx = \text{computer}$$

- k)  $\iint_S (x + z) dS$  where S is the first-octant portion of the cylinder  $y^2 + z^2 = 9$  between  $x = 0$  and  $x = 4$

$$z = \pm \sqrt{9 - y^2} \quad f_x = 0, \quad f_y = -\frac{y}{\sqrt{9 - y^2}}, \quad |f_x|^2 + |f_y|^2 = 1 + \frac{y^2}{9 - y^2} = \frac{9 - y^2 + y^2}{9 - y^2} = \frac{9}{9 - y^2}$$

$$\Rightarrow \int_0^4 \int_0^3 (x + \sqrt{9 - y^2}) \cdot \frac{3}{\sqrt{9 - y^2}} dy dx = \int_0^4 \int_0^3 (x + \sqrt{9 - y^2}) \frac{3}{\sqrt{9 - y^2}} dy dx = \text{computer}$$

7. For some of the following line integrals there may be short-cut you can use to simplify your computations (but justify your shortcut by quoting the appropriate theorem)

a)  $\int_C F \cdot dr$  where  $F(x, y) = \langle e^x \cos(y), -e^x \sin(y) \rangle$  and C is the curve  $r(t) = \langle 2\cos(t), 2\sin(t) \rangle, 0 \leq t \leq 2\pi$

closed curve so use Green:  $= \iint_D (-e^x \sin(y)) - (-e^x \cos(y)) dA = 0$

b)  $\int_C 2xyz dx + x^2 z dy + x^2 y dz$  where C is some smooth curve from (0,0,0) to (1,4,3)

one potential.  $f(x, y, z) = x^2 y z + C \Rightarrow = x^2 y z \Big|_{(0,0,0)}^{(1,4,3)} = 1 \cdot 4 \cdot 3 = 12$   
(checked)

c)  $\int_C F \cdot dr$  where  $F(x, y) = \langle y^3 + 1, 3xy^2 + 1 \rangle$  and C is the upper half of the unit circle, from (1,0) to (-1,0)

conservative with potential  $y^3 x + x + y + C \Big|_{(1,0)}^{(-1,0)} = -1 - 1 = -2$


d)  $\int_C F \cdot dr$  where  $F(x, y) = \langle y^3 x, 3xy^2 \rangle$  and C is the line segment from (-1,0) to (2,3).

No shortcut:  $r(t) = \langle -1, 0 \rangle + t \langle 3, 3 \rangle = \langle -1 + 3t, 3t \rangle, t \in [0, 1]$

$$\int_C y^3 x dx + 2xy^2 dy = \int_0^1 (3+3)^3 (-1+3t) \cdot 3 dt + 3(-1+3t)(3t)^2 \cdot 3 dt = \text{computer}$$

e)  $\int_C y^3 dx + (x^3 + 3xy^2) dy$  where C is the path from (0,0) to (1,1) along the graph of  $y = x^3$  and from (1,1) to (0,0) along the graph of  $y = x$ .

closed curve, so Green's theorem:  $= \iint_R (3x^2 + 3y) - 2y^2 dA = -3 \iint_R x^2 dA = -3 \int_0^1 \int_0^1 x^2 dy dx = \text{computer}$



8. Green's Theorem

- a) Use Green's theorem to find  $\int_C F \cdot dr$  where  $F(x, y) = \langle y^3, x^3 + 3xy^2 \rangle$  and C is the circle with radius 3, oriented counter-clockwise (You may need the double-angle formula for cos somewhere during your computations, or use Maple)

closed curve  $\oint_C \vec{F} \cdot d\vec{r} = \iint_{\text{Disk}} \underbrace{3x^2 + 3y^2}_{\substack{\text{div} \\ \text{of } \vec{F}}} - 3y^2 = \iint 3x^2 \, dA = \int \int 3r^2 \cos^2 \theta \, r \, dr \, d\theta =$  calculator

- b) Evaluate  $\iint_R dA$  where R is the ellipse  $\frac{x^2}{4} + \frac{y^2}{9} = 1$  by using a vector field  $F(x, y) = \langle -\frac{y}{2}, \frac{x}{2} \rangle$  and the boundary C of the ellipse R. Note that we did this in class, it is a very special application of Green's theorem.

$\vec{F} = \langle -\frac{y}{2}, \frac{x}{2} \rangle$

9. Evaluate the following integrals. You can use any theorem that's appropriate:

- c)  $\int_C 2xyz \, dx + x^2 z \, dy + x^2 y \, dz$  where C is a smooth curve from (0,0,0) to (1,4,3)

see previous examples

- d)  $\int_C y \, dx + 2x \, dy$  where C is the boundary of the square with vertices (0,0), (0,2), (2,0), and (2,2)


Green:  $\iint_{\text{square}} (2-1) \, dA = \text{area (square)} = \underline{4}$

- e)  $\oint_C xy^2 \, dx + x^2 y \, dy$ , where C is given by  $r(t) = \langle 4 \cos(t), 2 \sin(t) \rangle$ , t between 0 and 2 Pi.

Green, conservative vector field:  $= \underline{\underline{0}}$

- f)  $\int_C xy \, dx + x^2 \, dy$  where C is the boundary of the region between the graphs of  $y = x^2$  and  $y = x$ .

Green:  $= \iint_R (2x - x) \, dA = \int_0^1 \int_{x^2}^x x \, dx \, dy =$  calculator



10. Prove that if  $F(x, y, z) = \langle M(x, y, z), N(x, y, z), P(x, y, z) \rangle$  is any vector field where  $M, N, P$  are twice continuously differentiable then  $\text{div}(\text{curl}(F)) = 0$

just try it, it will work out

Use Green's Theorem to prove that integrals of a conservative vector fields over closed curves are zero (assuming that the closed curve encloses a simply connected region and all conditions of Green's theorem are satisfied).

extra credit...