

Math 2511: Calc III - Practice Exam 3

1. State the meaning or definitions of the following terms:

- a) vector field, conservative vector field, potential function of a vector field, volume, length of a curve, work, surface area, flux of a vector field, triple integral
 b) curl and divergence of a vector field F , gradient of a function

c) $\iint_R dA = \text{area}(R)$

$$\iint_R f(x, y) dA = \lim_{n \rightarrow \infty} \sum_{j=1}^n \sum_{i=1}^n f(x_i, y_j) \Delta x_i \Delta y_j = \int_a^b \int_c^d f(x, y) dy dx$$

$$\iiint_Q f(x, y, z) dV = \lim_{n \rightarrow \infty} \sum_{i=1}^n \sum_{j=1}^n \sum_{k=1}^n f(x_i, y_j, z_k) \Delta x_i \Delta y_j \Delta z_k =$$

d) $\int_C ds = \int_a^b \sqrt{(x')^2 + (y')^2} dt = \int_a^b \int_c^d \int_e^f f(x, y, z) dz dy dx$

$$\int_C f(x, y) ds = \int_a^b f(x(t), y(t)) \sqrt{(x')^2 + (y')^2} dt$$

$$\int_C f(x, y) dx = \int_a^b f(x(t), y(t)) x'(t) dt$$

$$\int_C f(x, y) dy = \int_a^b f(x(t), y(t)) y'(t) dt$$

e) $\int_C \vec{F} \cdot d\vec{r} = \int_C M dx + N dy + P dz$

$$\int_C M(x, y, z) dx + N(x, y, z) dy + P(x, y, z) dz$$

f) $\iint_S g(x, y, z) \cdot dS = \iint_R g(x, y, f(x, y)) \sqrt{1 + f_x^2 + f_y^2} dA, S: z = f(x, y)$

$$\iint_S \vec{F} \cdot \vec{n} dS = \iint_R -f_x M - f_y N + P dA, \text{ with } z = f(x, y)$$

g) What does it mean when a "line integral is independent of the path"?


If γ_1 and γ_2 are two paths, both starting at A and ending at B , then $\int_{\gamma_1} \vec{F} \cdot d\vec{r} = \int_{\gamma_2} \vec{F} \cdot d\vec{r}$

h) State the Fundamental Theorem of Line Integrals. Make sure to know when it applies, and when it helps

If C is a curve given by $\gamma: [a, b] \rightarrow \mathbb{R}^3$ and \vec{F} a conservative vector field with potential function f , then $\int_C \vec{F} \cdot d\vec{r} = f(\gamma(b)) - f(\gamma(a))$

i) State Green's Theorem. Make sure to know when it applies, and in what situation it helps.

If C is a closed curve and $\vec{F} = \langle M, N \rangle$ a 2D vector field such that all partials are continuous, then

$$\oint_C \vec{F} \cdot d\vec{r} = \iint_D \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dA$$


j) State Stoke's Theorem. Make sure to know when it applies, and in what situation it helps.

If S is a surface in \mathbb{R}^3 with a closed boundary curve C , all properly oriented then


$$\oint_C \vec{F} \cdot d\vec{r} = \iint_S \text{curl}(\vec{F}) \cdot \vec{n} \, dS$$

k) State Gauss' Theorem. Make sure to know when it applies, and in what situation it helps.

If S is a closed surface in \mathbb{R}^3 around Q then

$$\iint_S \vec{F} \cdot \vec{n} \, dS = \iiint_Q \text{div}(\vec{F}) \, dV$$


2. Below are four algebraic vector fields and four sketches of vector fields. Match them.



(1) $F(x, y) = \langle x, y \rangle$, (2) $F(x, y) = \langle -y, x \rangle$, (3) $F(x, y) = \langle x, 1 \rangle$, (4) $F(x, y) = \langle 1, y \rangle$

B
A
D
C

b) Below are two vector fields. Which one is clearly not conservative, and why?



so the 1st is not conservative

$$\oint_C \vec{F} \cdot d\vec{r} \neq 0$$

c) Say in the left vector field above you integrate over a straight line from $(0, -1)$ to $(1, 0)$. Is the integral positive, negative, or zero?

line and vectors are similar $\Rightarrow \int_C \vec{F} \cdot d\vec{r} > 0$


How about if you integrate from $(-2, 1)$ to $(2, 1)$?

line moves against vectors $\Rightarrow \int_C \vec{F} \cdot d\vec{r} < 0$

How about from (-2,-1) to (2,-1)?


$$\int_C \vec{F} \cdot d\vec{r} > 0$$

d) Below are two slices of a 3D surface submerged in a 3D vector field. Imagine the picture as extending outwards of the paper. The normal vector shown is determined by the orientation of the surface. Is the flux $\iint_S \vec{F} \cdot \vec{n} \, dS$ positive, negative, or zero, for each surface?




$$\vec{F} \cdot \vec{n} > 0$$

⇒ Flux positive



$$\vec{F} \cdot \vec{n} = 0$$

⇒ Flux zero



$$\vec{F} \cdot \vec{n} < 0$$

⇒ Flux negative

3. Are the following statements true or false:

a) If the divergence of a vector is zero, the vector field is conservative. ∇

b) If $F(x, y, z)$ is a conservative vector field then $\text{curl}(F) = 0$ ∇

c) If a line integral is independent of the path, then $\iint_S \vec{F} \cdot \vec{n} \, dS = 0$ for every path surface S ∇

d) If a vector field is conservative then $\int_C \vec{F} \cdot d\vec{r} = 0$ for every closed path C ∇

e) $\iint_R dA$ gives the surface area of the region R ∇

f) $\iint_R f(x, y) dA$ gives the volume of the region under the surface $f(x, y)$ and over R, if f is positive. ∇

g) $\iiint_Q dV$ gives the volume of Q ∇

h) Can you apply the Fundamental Theorem of line integrals for the function $f(x, y, z) = xy \sin(z) \cos(x^2 + y^2)$?
No (need vector field)

i) Can you apply the Fundamental Theorem of line integrals for the vector field $F(x, y) = \langle 6xy^2 - 3x^2, 6x^2y + 3y^2 - 7 \rangle$?
 $M_x = 12xy, M_y = 12xy$, so YES

j) Can you apply Green's theorem for a curve C, which is a straight line from (0,0,0) to (1,2,3)?
No (need closed curve)

k) Can you apply Green's theorem for a vector field $F = \langle xy, yz, zx \rangle$ and a closed surface S?

No, but Gauss's Theorem would work

l) Can you apply Gauss' theorem for a vector field $F = \langle xy, yz, zx \rangle$ and a surface S given by $z = f(x, y)$

No (need closed surface)

m) Can you apply Stoke's theorem for a 3D vector field and a surface S given by $z = f(x, y)$

Maybe (depends on surface)

4. Suppose that $F(x, y, z) = \langle x^3 y^2 z, x^2 z, x^2 y \rangle$ is some vector field. Find, if possible

a) $\text{div}(F), \quad \text{div} \{x^3 y^2 z, x^2 z, x^2 y\} = 3x^2 y^2 z$

$\text{curl}(F), \quad \text{curl} \{x^3 y^2 z, x^2 z, x^2 y\} = \{0, xy(x^2 y - 2), -2xz(x^2 y - 1)\}$

~~$\text{div}(\text{div}(F)),$~~

$\text{curl}(\text{curl}(F)), \quad \text{curl curl} \{x^3 y^2 z, x^2 z, x^2 y\} = \{-2x^3 z, 2z(3x^2 y - 1), y(3x^2 y - 2)\}$

$\text{div}(\text{curl}(F)), \quad 0$

~~$\text{curl}(\text{div}(F)),$~~

~~$\text{grad}(\text{curl}(F))$~~

$\text{grad}(\text{div}(F)) \quad \text{grad div} \{x^3 y^2 z, x^2 z, x^2 y\} = \{6xy^2 z, 6x^2 yz, 3x^2 y^2\}$

b) $\text{grad}., \text{div}.,$ and curl of the vector field if appropriate for $\langle x^2, y^2, z^2 \rangle$

grad d.u. apply

$\text{div}(F) = 2x + 2y + 2z, \quad \text{curl}(F) = \langle 0, 0, 0 \rangle$

c) $\text{grad}., \text{div}.,$ and curl of the vector field if appropriate for $\langle \cos(y) + y \cos(x), \sin(x) - x \sin(y), xyz \rangle$

grad d.u. apply

$\text{div}(F) = -y \sin(x) - x \cos(y) + xy, \quad \text{curl}(F) = \langle xz, -yz, 0 \rangle$

d) $\text{grad}., \text{div}.,$ and curl of the vector field if appropriate for $f(x, y, z) = z \ln(x^2 + y^2)$

$\text{grad}(F) = \left\langle \frac{z \cdot 2x}{x^2 + y^2}, \frac{z \cdot 2y}{x^2 + y^2}, \ln(x^2 + y^2) \right\rangle$ *curl and div d.u. apply*

5. Decide which of the following vector fields are conservative. If a vector is conservative, find its potential function

a) $F(x, y) = \langle 2xy, x^2 \rangle$

yes. $\underline{F = x^2 y + y x^2 + C}$

b) $F(x, y) = \langle e^x \cos(y), e^x \sin(y) \rangle$

No

c) $F(x, y, z) = \langle \sin(y), -x \cos y, 1 \rangle$

No (curl)

d) $F(x, y, z) = \langle 2xy, x^2 + z^2, 2zy \rangle$

Yes, $f = x^2 y + y z^2 + C$

e) $F(x, y) = \langle 6xy^2 - 3x^2, 6x^2 y + 3y^2 - 7 \rangle$

Yes, $f(x, y) = 3x^2 y^2 - x^3 + y^3 - 7y + C$

f) $F(x, y) = \langle -2y^3 \sin(2x), 3y^2(1 + \cos(2x)) \rangle$

Yes $M_x = -6y^3 \sin(2x), M_y = -6y^2 \sin(2x) \Rightarrow f = \cos(2x) y^3 + y^3 + C$

g) $F(x, y, z) = \langle 4xy + z, 2x^2 + 6y, 2z \rangle$

No; $\text{curl}(F) = \langle 0, 0, 0 \rangle$

h) $F(x, y, z) = \langle 4xy + z^2, 2x^2 + 6yz, 2xz \rangle$

No; $\text{curl}(4xy + z^2, 2x^2 + 6yz, 2xz) = \langle -6y, 0, 0 \rangle$

6. Evaluate the following integrals:

a) $\iint_R \cos(x^2) dA$ where R is the triangular region bounded by $y = 0$, $y = x$, and $x = 1$



$$\int_0^1 \int_0^x \cos(x^2) dy dx = \int_0^1 x \cos(x^2) dx = \frac{1}{2} \sin(1)$$

b) $\int_0^1 \int_1^{2y} x^2 y^3 dx dy$

$$\int_0^1 \int_1^{2y} x^2 y^3 dx dy = \frac{25}{84}$$

c) $\int_C ds$, where C is the curve given by $r(t) = \langle t^2, 1+t \rangle$, $0 \leq t \leq 2$ (you might want to use Maple at some point)

$$\int_0^2 \sqrt{(2t)^2 + 1} dt = \sqrt{17} + \frac{1}{4} \sinh^{-1}(4)$$

d) $\int x^2 y^3 dx$, where C is the curve given by $r(t) = \langle t^2, t^3 \rangle$, $0 \leq t \leq 2$

$$\int_0^2 (t^2)^2 (t^3)^3 \cdot \sqrt{4t^2 + 9t^4} dt = 2 \int_0^2 t^{14} dt = \frac{2}{15} 2^{15}$$

e) $\int_C x^2 - y + 3z ds$ where C is the circle $r(t) = \langle 2\cos(t), 2\sin(t) \rangle$, $0 \leq t \leq \pi$

$$\int_0^\pi ((2\cos(t))^2 - 2\sin(t) + 3) 2 dt = 10\pi - 8$$

f) $\int_C x^2 - y + 3z ds$ where C is a line segment given by $r(t) = \langle t, 2t, 3t \rangle$, $0 \leq t \leq 1$

$$\int_0^1 (t^2 - 2t + 3) \sqrt{1 + 2^2 + 3^2} dt = \frac{7\sqrt{14}}{3}$$

g) $\int_C F \cdot dr$ where $F(x, y) = \langle y, x^2 \rangle$ and C is the curve given by $r(t) = \langle 4-t, 4t-t^2 \rangle$, $0 \leq t \leq 3$

$$\int_0^3 ((4-t-t^2)(-1) + (4-t)^2(4-2t)) dt = \frac{69}{2}$$

h) $\int_C F \cdot dr$ where $F(x, y) = \langle yz, x^2, zy \rangle$ and C is the curve given by $r(t) = \langle 1-t, 3t, 2-t^2 \rangle$, $1 \leq t \leq 3$

$$\int_1^3 ((3t)(2-t^2)(-1) + (1-t)^2 3 + (2-t^2)(3t)(-2t)) dt = \frac{1152}{5}$$


i) $\int_C y dx + x^2 dy$ where C is a parabolic arc given by $r(t) = \langle t, 1-t^2 \rangle$, $-1 \leq t \leq 1$

$$\int_{-1}^1 ((1-t^2)1 + t^2(-2t)) dt = \frac{4}{3}$$

j) Find the surface integral $\iint_S x - 2y + z dS$, where S is the surface $z = 10 - 2x + 2y$ such that x is between 0 and 2 and y is between 0 and 4.

$$\int_0^4 \int_0^2 (x - 2y + (10 - 2x + 2y)) \sqrt{1 + 4 + 4} dx dy = 216$$

k) $\iint_S (x+z) dS$ where S is the first-octant portion of the cylinder $y^2 + z^2 = 9$ between $x=0$ and $x=4$



$$\int_0^4 \int_0^3 (x + \sqrt{9-y^2}) \sqrt{1 + 0 + \frac{y^2}{9-y^2}} dy dx = \underline{\underline{12(3+\pi)}}$$

7. For some of the following line integrals there may be short-cut you can use to simplify your computations (but justify your shortcut by quoting the appropriate theorem)

a) $\int_C F \cdot dr$ where $F(x, y) = \langle e^x \cos(y), -e^x \sin(y) \rangle$ and C is the curve $r(t) = \langle 2\cos(t), 2\sin(t) \rangle$, $0 \leq t \leq 2\pi$

C closed \Rightarrow Green's: $\int F \cdot dr = \iint -e^x \sin(y) + e^x \cos(y) dA = 0$
 (F is conservative)

b) $\int_C 2xyz dx + x^2 z dy + x^2 y dz$ where C is some smooth curve from (0,0,0) to (1,4,3)

$\int F \cdot dr$, $F = \langle 2xyz, x^2 z, x^2 y \rangle$ F has potential $f = x^2 y z$
 $= x^2 y z \Big|_{(0,0,0)}^{(1,4,3)} = 1 \cdot 4 \cdot 3 = \underline{\underline{12}}$

c) $\int_C F \cdot dr$ where $F(x, y) = \langle y^3 + 1, 3xy^2 + 1 \rangle$ and C is the upper half of the unit circle, from (1,0) to (-1,0)

$f = xy^3 + x + y$ is potential

$\Rightarrow \int F \cdot dr = xy^3 + x + y \Big|_{(1,0)}^{(-1,0)} = -1 - 1 = \underline{\underline{-2}}$

d) $\int_C F \cdot dr$ where $F(x, y) = \langle y^3 x, 3xy^2 \rangle$ and C is the line segment from (-1,0) to (2,3).

$\frac{\partial M}{\partial x} = 3y^2$, $\frac{\partial N}{\partial y} = 3y^2 x$ Not conservative $r(t) = \langle -1, 0 \rangle + t \langle 3, 3 \rangle = \langle -1 + 3t, 3t \rangle$
 $\int F \cdot dr = \int_0^1 (3t)^3 \cdot (3t) + \int_0^1 (3t+1)(3t)^2 (3) dt = \underline{\underline{45}}$

e) $\int_C y^3 dx + (x^3 + 3xy^2) dy$ where C is the path from (0,0) to (1,1) along the graph of $y = x^3$ and from (1,1) to (0,0) along the graph of $y = x$.



closed curve \Rightarrow Green's: $\int F \cdot dr = \iint (3x^2 + 3y^2 - 3y^2) dA = \iint 3x^2 dy dx = \int_0^1 \int_0^x 3x^2 dy dx = \underline{\underline{\frac{3}{4}}}$

8. Green's Theorem

- a) Use Green's theorem to find $\int_C \vec{F} \cdot d\vec{r}$ where $F(x, y) = \langle y^3, x^3 + 3xy^2 \rangle$ and C is the circle with radius 3, oriented counter-clockwise (You may need the double-angle formula for cos somewhere during your computations, or use Maple)

$$\int_C \vec{F} \cdot d\vec{r} = \iint_D (3x^2 + 3y^2 - 3y^2) dA = \iint_D 3x^2 dx dy =$$

$$\stackrel{\text{circle}}{=} \int_0^{2\pi} \int_0^3 3r^2 \cos^2 \theta + dr d\theta = \underline{\underline{\frac{3\pi}{4}}}$$

- b) Evaluate $\iint_R dA$ where R is the ellipse $\frac{x^2}{4} + \frac{y^2}{9} = 1$ by using a vector field $F(x, y) = \langle -\frac{y}{2}, \frac{x}{2} \rangle$ and the boundary C of the ellipse R. Note that we did this in class, it is a very special application of Green's theorem.

- c) Find the surface integral $\iint_S x - 2y + z dS$, where S is the surface $z = 10 - 2x + 2y$ such that x is between 0 and 2 and y is between 0 and 4.

$$\iint_R x - 2y + (10 - 2x + 2y) \sqrt{9} dA = \int_0^2 \int_0^4 10 - x dy dx = \underline{\underline{72}}$$

- d) Evaluate the flux integral $\iint_S \vec{F} \cdot \vec{n} dS$ where $F(x, y, z) = \langle x, y, z \rangle$ and S is $x^2 + y^2 + z^2 = 4$


Gauss Thm: Flux = $\iiint_{\text{sphere}} \text{div}(\vec{F}) dV =] \cdot \text{Volume (sphere)}$

$$= 3 \cdot \frac{4}{3} \pi 2^3 = \underline{\underline{2\pi}}$$

e) Evaluate $\int_C \vec{F} d\vec{r}$ where $F(x, y, z) = \langle z^2, x^2, y^2 \rangle$ and C is the boundary of the surface S given by $z = 4 - x^2 - y^2$ and $z \geq 0$, oriented counter-clockwise.

Stoke's $\oint_C \vec{F} \cdot d\vec{s} = \iiint_V \text{div}(\vec{F}) dV = 0$


f) Evaluate $\int_C \vec{F} d\vec{r}$ where $F(x, y, z) = \langle -y^2, x, z^2 \rangle$ and C is the curve bounding the ellipse S consisting of the intersection of the plane $y + z = 2$ and the cylinder $x^2 + y^2 = 1$



$$\int_C \vec{F} \cdot d\vec{r} = \iint_S \text{curl}(\vec{F}) \cdot \vec{n} dA = \iint_D \langle 0, 0, 2y+1 \rangle \cdot \langle 0, 1, 1 \rangle dA$$

$$= \iint_{\text{disk}} 2y+1 dA = \int_0^{2\pi} \int_0^1 (2r \sin(\theta) + 1) r dr d\theta = \pi$$

g) Evaluate $\iint_S \text{curl}(\vec{F}) \cdot \vec{n} dS$ where $F(x, y, z) = \langle xz, yz, xy \rangle$ and S is the part of the sphere $x^2 + y^2 + z^2 = 4$ that lies inside the cylinder $x^2 + y^2 = 1$ above the xy-plane.



Use Stoke's Theorem: $\iint_S \text{curl}(\vec{F}) \cdot \vec{n} dS = \int_C \vec{F} \cdot d\vec{r} = \int_C xz dx + yz dy + xy dz = 0$

Use Stoke's Theorem: $\iint_S \text{curl}(\vec{F}) \cdot \vec{n} dS = \int_C \vec{F} \cdot d\vec{r} = \int_C xz dx + yz dy + xy dz = 0$

$C: \vec{r}(t) = \langle \cos(t), \sin(t), k \rangle$

9. Evaluate the following integrals. You can use any theorem that's appropriate:

h) $\int_C 2xyz dx + x^2 z dy + x^2 y dz$ where C is a smooth curve from (0,0,0) to (1,4,3)

$f = x^2 y z$ is potential \Rightarrow

$$\int_C \vec{F} \cdot d\vec{r} = x^2 y z \Big|_{(0,0,0)}^{(1,4,3)} = \underline{\underline{12}}$$

$$= \int_C k \cos(t)(-\sin(t)) + k \sin(t) \cos(t) dz = 0 \quad \underline{\underline{0}}$$


i) $\int_C y dx + 2x dy$ where C is the boundary of the square with vertices (0,0), (0,2), (2,0), and (2,2)

closed curve, Green: $\iint_{\text{square}} 2-1 dA = \text{area}(\text{square}) = \underline{\underline{4}}$

j) $\int_C xy^2 dx + x^2 y dy$, where C is given by $r(t) = \langle 4 \cos(t), 2 \sin(t) \rangle$, t between 0 and 2π . ellipse

closed curve \rightarrow Green: $\iint_D 2xy - 2yx \, dA = \underline{\underline{0}}$

k) $\int_C xy dx + x^2 dy$ where C is the boundary of the region between the graphs of $y = x^2$ and $y = x$.



Green: $\iint_D 2x - x \, dA = \int_0^1 \int_{x^2}^x x \, dy dx = \underline{\underline{\frac{2}{15}}}$

10. Prove that if $F(x, y, z) = \langle M(x, y, z), N(x, y, z), P(x, y, z) \rangle$ is any vector field where M, N, P are twice continuously differentiable then $\text{div}(\text{curl}(F)) = 0$

$$\text{curl}(F) = \begin{vmatrix} i & j & k \\ \partial_x & \partial_y & \partial_z \\ M & N & P \end{vmatrix} = \langle P_y - N_z, -(P_x - M_z), N_x - M_y \rangle$$

$$\begin{aligned} \text{div}(\text{curl}(F)) &= (P_y - N_z)_x - (P_x - M_z)_y + (N_x - M_y)_z \\ &= \cancel{P_{yx}} - \cancel{N_{zx}} - \cancel{P_{xy}} + \cancel{M_{zy}} + \cancel{N_{xz}} - \cancel{M_{yz}} = 0 \quad \# \end{aligned}$$

Use Green's Theorem to prove that integrals of a conservative vector fields over closed curves are zero (assuming that the closed curve encloses a simply connected region and all conditions of Green's theorem are satisfied).

$$C \text{ closed} \rightarrow \oint_C F \, dr = \iint_D \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \, dA$$

$$F \text{ conservative} \Rightarrow \frac{\partial N}{\partial x} = \frac{\partial M}{\partial y} \Rightarrow \oint_C F \, dr = \iint_D 0 \, dA = 0 \quad \#$$