

The Legalization of Recreational Marijuana: How Likely Is the Worst-Case Scenario?

D. Mark Anderson and Daniel I. Rees

INTRODUCTION

Last fall, voters in Colorado and Washington approved measures legalizing the recreational use of marijuana. In the near future, residents of these states who are 21 years of age and older will be able to purchase marijuana at retail stores (Donlan, 2013). Although it can be challenging to predict future behavior, Mark Kleiman, a prominent drug policy expert, described what might be characterized as the worst-case scenario. According to Kleiman, this scenario would involve three elements: more heavy drinking, “carnage on our highways,” and a “massive” increase in the use of marijuana by minors (Livingston, 2013).

Below, we discuss the likely effects of legalizing marijuana for recreational use on alcohol consumption, traffic fatalities, substance use among high school students, and other outcomes of interest to policymakers and the public. Our discussion draws heavily on studies that have examined the legalization of medical marijuana. These studies are relevant because, in states such as California, Colorado, Oregon, and Washington, the legalization of marijuana for medicinal purposes approaches de facto legalization of marijuana for recreational purposes.

One of the key unknowns in the debate over legalization concerns the relationship between alcohol and marijuana use. Researchers have attempted to produce causal estimates of this relationship by exploiting cross-sectional policy and price variation (Pacula, 1998; Williams et al., 2004). We note that these estimates could easily be spurious and that more reliable estimates based on clearly defined natural experiments show that alcohol and marijuana are substitutes. Because the social costs associated with the consumption of alcohol clearly outweigh those associated with the consumption of marijuana, we conclude that legalizing the recreational use of marijuana is likely to improve public health, although plenty of unanswered questions remain.

BACKGROUND

In 1996, California voters approved the Compassionate Use Act that removed criminal penalties for using, possessing, and cultivating medical marijuana. Under this act, doctors are allowed to recommend the use of marijuana “in the treatment of cancer, anorexia, AIDS, chronic pain, spasticity, glaucoma, arthritis, migraine, or any other illness for which marijuana provides relief,” and patients are allowed to designate a caregiver, who can obtain or grow marijuana on their behalf.¹

The Compassionate Use Act affords growers and distributors some protection against prosecution and allows patients to buy marijuana without fear of being

¹ The full text of the Compassionate Use Act of 1996 is available at <http://www.cdph.ca.gov/programs/mmp/pages/compassionateuseact.aspx>.

arrested or fined. Because it is prohibitively expensive to ensure that all medical marijuana ends up in the hands of patients, a substantial portion of the California medicinal crop is diverted to the recreational market (Montgomery, 2010; Nagourney, 2012). Since 1996, 20 additional states have passed medical marijuana laws (although in an effort to prevent diversion to the recreational market, a number of these states limit caregivers to one patient, prohibit home cultivation, or prohibit dispensaries). By examining pre- and post-legalization data from these states, we can make predictions about what will happen in Colorado and Washington.

THE EFFECT OF LEGALIZATION ON PRICE

The legalization of medical marijuana can be thought of as increasing both the supply of and the demand for marijuana (Pacula et al., 2010). Although the effect of legalizing medical marijuana on consumption is unambiguously positive, its effect on price could be either positive or negative.

Anderson, Hansen, and Rees (2013) collected price data from back issues of *High Times* magazine for the period 1990 to 2011 to gauge the impact of legalizing medical marijuana on the marijuana market. They found that legalization was associated with a 10 to 26 percent decrease in the price of high-quality marijuana, suggesting the supply response to legalizing medical marijuana is larger than the demand response. These authors also compared *High Times* prices to prices advertised by dispensaries in Arizona, California, Colorado, Michigan, Nevada, Oregon, and Washington. The prices were similar, suggesting that there is substantial overlap between the medicinal and recreational markets in these states.

Both Colorado and Washington have legalized the production and commercial distribution of marijuana for recreational purposes, but these activities are still prohibited under federal law. As a consequence, large-scale farming of marijuana with tractors and unskilled workers is not likely to occur (Caulkins et al., 2012, pp. 192–193). Instead, most production will continue to take place under grow lights inside relatively small facilities. Because the production of medical marijuana under these conditions is already widespread in Colorado and Washington, price may not fall much further than it already has. If a nonmedical marijuana state such as New York were to legalize the recreational use of marijuana without first taking the intermediate step of legalizing medicinal use, the price of marijuana would, in all likelihood, fall substantially. The results of Anderson, Hansen, and Rees (2013) suggest that, under these circumstances, the price of marijuana could fall by as much as 40 percent after four to five years.

MARIJUANA USE

The National Survey on Drug Use and Health (NSDUH) is the best source of information on marijuana consumption by adults living in the United States. However, the NSDUH does not typically provide individual-level data with state identifiers to researchers, and did not publish state-level estimates of marijuana use prior to 1999. Because five states (including California, Oregon, and Washington) legalized medical marijuana during the period 1996 to 1999, the NSDUH is of limited value when trying to estimate the effect of legalization on the use of marijuana.

As an alternative to using NSDUH data, we examined the effect of legalizing medical marijuana on the amount of marijuana eradicated under the Domestic Cannabis Eradication/Suppression Program during the period 1990 to 2010. Standard difference-in-differences estimates based on data at the state-year level are reported in Table 1.

Table 1. The legalization of medical marijuana and Drug Enforcement Administration (DEA) eradication efforts.

	ln(total plants eradicated)		ln(outdoor plants eradicated)		ln(indoor plants eradicated)	
	(1)	(2)	(3)	(4)	(5)	(6)
Legalization of medical marijuana	1.08*** (0.240)	0.113 (0.212)	1.289** (0.389)	-0.045 (0.220)	0.415* (0.240)	0.608* (0.311)
<i>N</i>	1040	1040	963	963	972	972
<i>R</i> ²	0.859	0.899	0.838	0.896	0.823	0.850
State and year FEs	Yes	Yes	Yes	Yes	Yes	Yes
State-specific trends	No	Yes	No	Yes	No	Yes

*Statistically significant at the 10 percent level; **at the 5 percent level; ***at the 1percent level.

Notes: Based on state-level data produced by the Domestic Cannabis Eradication/Suppression Program for the period 1990 to 2010 available at <http://www.drugscience.org/Archive/DCESP/> and http://www.albany.edu/sourcebook/ind/DRUG_ENFORCEMENT_ADMINISTRATION.Seizures_and_removals.1.html. Each odd-numbered column represents the results from a separate OLS regression of the form:

$$\ln(\text{Plants eradicated}_{st}) = \beta_0 + \beta_1 \text{Legalization}_{st} + X_{st} \beta_2 + v_s + w_t + \varepsilon_{st},$$

where *s* indexes state and *t* indexes year. Each even-numbered column represents the results of a separate OLS regression that also includes state-specific linear time trends. The covariates composing the vector X_{st} are population, mean age, income, and the unemployment rate. Estimates are weighted using state populations. Standard errors, corrected for clustering at the state level, are in parentheses. Fourteen states implemented a medical marijuana law during the period 1990 to 2010.

Without state-specific linear time trends, legalizing medical marijuana is associated with an almost 200 percent increase in the number of marijuana plants eradicated under the Domestic Cannabis Eradication/Suppression Program ($e^{1.08} - 1 = 1.94$). This estimate, however, becomes much smaller and insignificant when state-specific linear time trends are added to the model. Without state-specific linear time trends, legalization is associated with a 51.4 percent increase in the number of indoor plants eradicated. When state-specific linear time trends are included, this estimate actually becomes larger: Legalization is associated with an 83.7 percent increase in the number of indoor plants eradicated.

These results are consistent with the hypothesis that legalizing medical marijuana leads to a substantial increase in the supply of marijuana grown indoors, but could reflect greater effort on the part of the Drug Enforcement Administration (DEA), the agency tasked with running the Domestic Cannabis Eradication/Suppression Program. Unfortunately, the DEA provides almost no information on the intensity of their eradication efforts. However, we do know that state officials and lawmakers from medical marijuana states have put pressure on the DEA to scale back its efforts (Aiello, 2011; Graves, 2012; Ingold, 2012; Slevin, 2010). In addition, if the post-legalization increase in indoor plants eradicated were due to a ramping up of effort, price should have increased. Instead, the price of high-quality marijuana fell after legalization (Anderson, Hansen, & Rees, 2013), suggesting that any increase in effort did not keep up with supply. If the DEA allows Colorado and Washington growers to expand their operations, a further decrease in price and a corresponding increase in consumption can be expected. By how much might consumption increase? The answer to this question depends, in part, on the price elasticity of demand for marijuana. A number of studies have attempted to estimate this elasticity (DeSimone & Farrelly, 2003; Nisbet & Vakil, 1972; Pacula et al., 2001; Williams et al., 2004), but

none exploited a clearly defined natural experiment. As a consequence, very little is known about how marijuana consumption would respond to further reductions in price.

ALCOHOL USE

Pacula (1998) and Williams et al. (2004) found evidence of a negative relationship between beer taxes and marijuana use, suggesting that beer and marijuana are complements. However, both of these studies relied on cross-sectional variation in state beer taxes, which could reflect difficult-to-observe factors, such as attitudes towards substance use. By way of an example, Utah taxes beer at five times the Colorado rate for historical and cultural reasons; these same reasons probably explain why marijuana use in Utah is so low as compared to Colorado.²

Exploiting within-state variation in the price of beer over time, Farrelly et al. (1999) found evidence of complementarity between alcohol and marijuana among teens, but not among young adults. Yörük and Yörük (2011) used data from the National Longitudinal Survey of Youth 1997 (NLSY97) and a regression discontinuity design to examine the effect of the minimum legal drinking age (MLDA) on marijuana use. They also found evidence of complementarity, but inadvertently conditioned on having used marijuana at least once since the last interview. When Crost and Rees (2013) applied Yörük and Yörük's (2011) research design to the NLSY97 data without conditioning on having used marijuana since the last interview, they found no evidence that alcohol and marijuana were complements.

Studies based on clearly defined natural experiments generally support the hypothesis that marijuana and alcohol are substitutes. For instance, DiNardo and Lemieux (2001) found that increasing the MLDA from 18 to 21 encourages marijuana use. Using data from the NSDUH and a regression discontinuity design, Crost and Guerrero (2012) found a sharp decrease in marijuana use at 21 years of age, suggesting that young adults treat alcohol and marijuana as substitutes. Finally, Anderson, Hansen, and Rees (2013) examined the relationship between legalizing medical marijuana and drinking using data from the Behavioral Risk Factor Surveillance System. These authors found that legalization was associated with reductions in heavy drinking especially among 18- through 29-year olds. In addition, they found that legalization was associated with an almost 5 percent decrease in beer sales, the alcoholic beverage of choice among young adults (Jones, 2008).

The results of DiNardo and Lemieux (2001), Crost and Guerrero (2012), and Anderson, Hansen, and Rees (2013) suggest that, as marijuana becomes more available, young adults in Colorado and Washington will respond by drinking less, not more. If nonmedical marijuana states legalize the use of recreational marijuana, they should also experience reductions in drinking with the accompanying public-health benefits.

TRAFFIC FATALITIES

Reducing traffic injuries and fatalities is potentially one of the most important public-health benefits from legalizing the use of recreational marijuana. Tetrahydrocannabinol (THC), the principal psychoactive component of marijuana, impairs

² According to the Tax Foundation, Utah, taxes beer at 41 cents per gallon, while Colorado taxes beer at 8 cents per gallon. According to data from the 2009 and 2010 NSDUH, approximately 3 percent of Utah residents ages 12 and over used marijuana in the past month. In comparison, approximately 11 percent of Colorado residents used marijuana in the past month (National Survey on Drug Use and Health, 2012).

driving-related functions (Kelly, Darke, & Ross, 2004), but there is evidence that drivers under the influence of THC compensate for these impairments. For instance, they tend to drive slower and take fewer risks (Robbe & O'Hanlon, 1993; Sewell, Poling, & Sofuoglu, 2009). In contrast, drivers under the influence of alcohol tend to drive faster and take more risks (Burian, Liguori, & Robinson, 2002; Marczynski, Harrison, & Fillmore, 2008; Ronen et al., 2008). While driving under the influence of marijuana is associated with a twofold increase in the risk of being involved in a collision (Asbridge, Hayden, & Cartwright, 2012), driving with a blood alcohol concentration (BAC) of 0.08 or greater is associated with a 4- to 27-fold increase in this same risk (Peck et al., 2008).

Driving under the combined influence of alcohol and marijuana is especially dangerous (Sewell, Poling, & Sofuoglu, 2009). Therefore, if young adults viewed alcohol and marijuana as complements, legalizing the recreational use of marijuana could seriously jeopardize roadway safety. Fortunately, as noted above, studies based on clearly defined natural experiments suggest that young adults, a group responsible for a disproportionate share of traffic accidents and fatalities (Eustace & Wei, 2010), typically substitute marijuana in place of alcohol.

Using data from the Fatality Analysis Reporting System (FARS) for the period 1990 to 2010, Anderson, Hansen, and Rees (2013) examined the effect of legalizing medical marijuana on traffic fatalities. They found that legalizing medical marijuana was associated with a 13 percent decrease in fatalities involving alcohol. It is important to note, however, that their results do not necessarily imply that driving under the influence of marijuana is safer than driving under the influence of alcohol. Because marijuana is not typically consumed in public venues, such as restaurants and bars, their results may reflect fewer impaired drivers on the road.³

Finally, the Washington law that legalized recreational marijuana also established a THC limit for drivers of 5 ng/ml of blood.⁴ A strict THC limit above which drivers are automatically considered impaired may, in the future, be viewed by the public and policymakers as a necessary complement to legalizing recreational marijuana. However, there is little evidence that, as currently implemented, such limits actually improve roadway safety (Anderson & Rees, 2012). Given this lack of evidence, policymakers might consider lowering the BAC limit. This could deter drunk driving (Dee, 2001; Eisenberg, 2003) and come with the added benefit of discouraging driving under the combined influence of alcohol and marijuana. Alternatively, a substantial increase in alcohol excise taxes could help to discourage driving under the combined influence of alcohol and marijuana. The current excise tax on liquor sold in Colorado is 60.26 cents/l, which represents roughly 3 percent of the retail price of Jim Beam Whiskey purchased by the bottle.⁵ In comparison, Colorado is set to impose a 15 percent excise tax and a 10 percent special sales tax on marijuana sales (Ingold, 2013b). Washington is considering taxing producers, sellers, and buyers at a total rate of 75 percent (Donlan, 2013).

³ A small number of quasi-private bars and clubs catering to marijuana users have begun operating in Colorado and Washington since the legalization of recreational marijuana (Johnson, 2013). If these establishments become popular, this could lead to more traffic fatalities involving drivers under the influence of marijuana.

⁴ The Colorado legislature voted to establish a similar threshold in May 2013 (Ingold, 2013a).

⁵ The retail price of Jim Beam Whiskey was obtained from Argonauts Liquor in Denver (<http://www.argonautliquor.com>), Davidson Liquors in Highlands Ranch (<http://davidsonliquors.com>), and Cheers Liquor Mart in Colorado Springs (<http://www.cheersliquormart.com>). Jim Beam is the best-selling brand of whiskey in the United States.




Figure 1. Marijuana Use Among Los Angeles High School Students, 2001 to 2011.

MARIJUANA USE BY TEENAGERS

Policymakers and the public are especially concerned that legalization of recreational marijuana will encourage substance use among teenagers. This concern is bolstered by the fact that Denver teenagers receiving treatment for substance abuse have reported obtaining marijuana from medical marijuana patients (Salomonsen-Sautel et al., 2012; Thurstone, Lieberman, & Schmiede, 2011). The interesting question, however, is not whether diversion to teenagers will occur with the legalization of recreational marijuana (it will). The more interesting (and policy-relevant) question is whether marijuana use by teenagers will increase when “pot shops” open throughout Colorado and Washington.

Using data from national and state Youth Risk Behavior Surveys (YRBS) for the period 1993 to 2011, Anderson, Hansen, and Rees (2012) found little evidence of a relationship between legalizing medical marijuana and the use of marijuana among high school students.⁶ Critics of Anderson, Hansen, and Rees (2012) have argued that youth access to medical marijuana did not appreciably increase until medical marijuana dispensaries became widespread (Ferner, 2012). In an effort to explore the role of dispensaries, we turned to local YRBS data for the period 2001 to 2011.⁷ Specifically, we compared marijuana use among Los Angeles high school students to marijuana use among Chicago, Dallas, Boston, and Miami-Dade County high school students (Figure 1). The first dispensaries began operating in Los Angeles prior to the passage of the Compassionate Use Act (Curtius & Yates, 1996), but the dispensary boom did not occur until after 2004 (Jacobson et al., 2011). From 2005 to 2010,

⁶ See also Harper, Strumpf, and Kaufman (2012) and Lynne-Landsman, Livingston, and Wagenaar (2013).

⁷ These data are available at <http://apps.nccd.cdc.gov/youthonline/App/Default.aspx?SID=HS>.

the number of dispensaries operating in Los Angeles surged from only a handful to more than 600 (Jacobson et al., 2011).⁸ There were no legal medical marijuana dispensaries operating in Chicago, Dallas, Boston, or Miami-Dade County during the period 2001 to 2011.

Figure 1 provides little evidence that marijuana use among Los Angeles high school students increased in the mid-2000s, perhaps because dispensaries could sell to adults with some assurance of not being shut down, while selling marijuana to a minor was still a risky proposition even after the passage of the Compassionate Use Act.⁹ Marijuana use among Los Angeles high school students actually declined from 2007 to 2009 and then increased from 2009 to 2011. However, there were comparably sized increases in marijuana use among Chicago, Dallas, and Boston high school students from 2009 to 2011.

CRIME AND OTHER PUBLIC-HEALTH OUTCOMES

There is evidence of a link between alcohol abuse and violent crime, including domestic violence (Carpenter & Dobkin, 2010; Markowitz, 2000, 2005; Markowitz & Grossman, 1998). Therefore, if the legalization of recreational marijuana leads to reduced alcohol consumption, we expect violent crime to fall. It is also possible that nonviolent crime will fall as policing resources are freed up and reallocated (Miron & Zwiebel, 1995). Adda, McConnell, and Rasul (2011) found that nondrug crime went down after Lambeth, a borough of London, temporarily suspended arrests for marijuana possession.

Other potential public-health benefits from legalizing recreational marijuana could include a reduction in suicides (Anderson, Rees, & Sabia, in press), increased product quality, and fewer emergency room episodes resulting from the use of marijuana laced with chemicals, such as phencyclidine (PCP) and embalming fluid. Quality oversight by the government is absent for illegal drugs, and repeated interactions between buyers and sellers in a prohibited market is, in all likelihood, only a partial solution (Miron & Zwiebel, 1995).

Finally, it is possible that the legalization of recreational marijuana will affect the demand for hard drugs, such as cocaine and heroin. Previous studies have documented a positive association between the use of marijuana and hard drugs (Bretteville-Jensen, Melberg, & Jones, 2008; DeSimone, 1998; Fergusson, Boden, & Horwood, 2006; Melberg, Jones, & Bretteville-Jensen, 2010; van Ours, 2003), but these studies were not based on clearly defined natural experiments. Two exceptions are Kelly and Rasul (2012) and Model (1993). Kelly and Rasul found that hospital admissions for hard drugs increased after the borough of Lambeth suspended arrests for marijuana possession. In contrast, Model, who used U.S. data, found that marijuana decriminalization led to fewer emergency room episodes involving drugs other than marijuana.

CONCLUSION

Kleiman's worst-case scenario is possible, but not likely. Based on existing empirical evidence, we expect that the legalization of recreational marijuana in Colorado and

⁸ In June of 2010, the City of Los Angeles ordered 439 (or approximately 70 percent) of its medical marijuana dispensaries to close (Jacobson et al., 2011, p. 8).

⁹ Federal authorities in Los Angeles have focused their attention on medical marijuana dispensaries that violate state law by, for instance, selling to minors (Meyer & Glover, 2009). Medical marijuana patients under the age of 18 must be accompanied by a parent or legal caregiver when visiting a dispensary (Ellison, 2009).

Washington will lead to increased marijuana consumption coupled with decreased alcohol consumption. As a consequence, these states will experience a reduction in the social harms resulting from alcohol use. While it is more than likely that marijuana produced by state-sanctioned growers will end up in the hands of minors, we predict that overall youth consumption will remain stable. On net, we predict the public-health benefits of legalization to be positive.

D. MARK ANDERSON is Assistant Professor at Department of Agricultural Economics and Economics, Montana State University, P.O. Box 172920, Bozeman, MT 59717-2920.

DANIEL I. REES is Professor at Department of Economics, University of Colorado Denver, CB 181, Denver, CO 80217-3364.

ACKNOWLEDGMENTS

The authors would like to thank Benjamin Crost and Jeffrey Miron for their comments and suggestions.

REFERENCES

- Adda, J., McConnell, B., & Rasul, I. (2011). Crime and the depenalization of cannabis possession: Evidence from a policing experiment. Working paper, Department of Economics, University College London. Retrieved September 24, 2013, from <http://www.ucl.ac.uk/~uctpimr/research/>.
- Aiello, D. (2011). Democrats call on feds to back off attacks on medical marijuana. California Progress Report. Retrieved September 24, 2013, from <http://www.californiaprogressreport.com/site/democrats-call-feds-back-attacks-medical-marijuana>.
- Anderson, D. M., & Rees, D. I. (2012). Per se drugged driving laws and traffic fatalities. IZA Discussion Paper No. 7048. Bonn, Germany: Institute for the Study of Labor.
- Anderson, D. M., Hansen, B., & Rees, D. I. (2012). Medical marijuana laws and teen marijuana use. Working Paper, Department of Agricultural Economics and Economics, Montana State University. Retrieved September 24, 2013, from <http://dmarkanderson.com/research>.
- Anderson, D. M., Hansen, B., & Rees, D. I. (2013). Medical marijuana laws, traffic fatalities, and alcohol consumption. *Journal of Law and Economics*, 56, 333–369.
- Anderson, D.M, Rees, D. I., & Sabia, J. J. (in press). Medical marijuana laws and suicides. *American Journal of Public Health*.
- Asbridge, M., Hayden, J., & Cartwright, J. (2012). Acute cannabis consumption and motor vehicle collision risk: Systematic review of observational studies and meta-analysis. *British Medical Journal*, 344, e536.
- Bretteville-Jensen, A. L., Melberg, H. O., & Jones, A. M. (2008). Sequential patterns of drug use initiation—Can we believe in the gateway theory? *B.E. Journal of Economic Analysis and Policy*, 8, Article 1. Retrieved September 24, 2013, from <http://www.degruyter.com/view/j/bejeap.2008.8.2/bejeap.2008.8.2.1846/bejeap.2008.8.2.1846.xml?format=INT>.
- Burian, S., Liguori, A., & Robinson, J. (2002). Effects of alcohol on risk-taking during simulated driving. *Human Psychopharmacology*, 17, 141–150.
- Carpenter, C., & Dobkin, C. (2010). The drinking age, alcohol consumption, and crime. Working Paper. Irvine, CA: Paul Merage School of Business, University of California-Irvine.
- Caulkins, J. P., Hawken, A., Kilmer, B., & Kleiman, M. R. A. (2012). *Marijuana legalization: What everyone needs to know*. New York, NY: Oxford University Press.
- Crost, B., & Guerrero, S. (2012). The effect of alcohol availability on marijuana use: Evidence from the minimum legal drinking age. *Journal of Health Economics*, 31, 112–121.

- Crost, B., & Rees, D. I. (2013). The minimum legal drinking age and marijuana use: New estimates from the NLSY97. *Journal of Health Economics*, 32, 474–476.
- Curtius, M., & Yates, N. (1996). Cannabis buyers' club founder arrested. *Los Angeles Times*. Retrieved September 24, 2013, from http://articles.latimes.com/1996-10-12/news/mn-53188_1_cannabis-buyers-club.
- Dee, T. (2001). Does setting limits save lives? The case of 0.08 BAC laws. *Journal of Policy Analysis and Management*, 20, 111–128.
- DeSimone, J. (1998). Is marijuana a gateway drug? *Eastern Economic Journal*, 24, 149–164.
- DeSimone, J., & Farrelly, M. C. (2003). Price and enforcement effects on cocaine and marijuana demand. *Economic Inquiry*, 41, 98–115.
- DiNardo, J., & Lemieux, T. (2001). Alcohol, marijuana, and American youth: The unintended consequences of government regulation. *Journal of Health Economics*, 20, 991–1010.
- Donlan, T. G. (2013). Should pot be legal? *Barron's*. Retrieved September 24, 2013, from http://online.barrons.com/article/SB50001424052748704509304578511261557343002.html#articleTabs_article%3D1.
- Eisenberg, D. (2003). Evaluating the effectiveness of policies related to drunk driving. *Journal of Policy Analysis and Management*, 22, 249–274.
- Ellison, K. (2009). Medical marijuana: No longer just for adults. *The New York Times*. Retrieved September 24, 2013, from <http://www.nytimes.com/2009/11/22/health/22sfmedical.html>.
- Eustace, D., & Wei, H. (2010). The role of driver age and gender in motor vehicle fatal crashes. *Journal of Transportation Safety and Security*, 2, 28–44.
- Farrelly, M. C., Bray, J. W., Zarkin, G. A., Wendling, B. W., & Pacula, R. L. (1999). The effects of prices and policies on the demand for marijuana: Evidence from the National Household Surveys on Drug Abuse. NBER Working Paper No. 6940. Cambridge, MA: The National Bureau of Economic Research.
- Fergusson, D. M., Boden, J. M., & Horwood, L. J. (2006). Cannabis use and other illicit drug use: Testing the cannabis gateway hypothesis. *Addiction*, 101, 556–569.
- Ferner, M. (2012). Marijuana usage down among Colorado teens, up nationally: Study shows. *Huffington Post*. Retrieved September 24, 2013, from http://www.huffingtonpost.com/2012/09/07/marijuana-usage-down-in-t_n_1865095.html.
- Graves, L. (2012). Lawmakers in 5 states tell feds to back off medical marijuana. *Huffington Post*. Retrieved September 24, 2013, from http://www.huffingtonpost.com/2012/04/02/lawmakers-in-5-states-tell-feds-medical-marijuana_n_1397811.html.
- Harper, S., Strumpf, E., & Kaufman, J. (2012). Do medical marijuana laws increase marijuana use? Replication study and extension. *Annals of Epidemiology*, 22, 207–212.
- Ingold, J. (2012). Boulder DA asks feds to back off on medical marijuana dispensaries. *The Denver Post*. Retrieved September 24, 2013, from http://www.denverpost.com/breakingnews/ci_20172102/boulder-da-asks-feds-back-off-medical-marijuana.
- Ingold, J. (2013a). Colorado Legislature gives final OK to marijuana driving limit. *The Denver Post*. Retrieved September 24, 2013, from http://www.denverpost.com/breakingnews/ci_23189484/colorado-legislature-gives-final-ok-marijuana-driving-limit.
- Ingold, J. (2013b). Colorado Legislature gives final approval to historic marijuana bills. *The Denver Post*. Retrieved September 24, 2013, from http://www.denverpost.com/breakingnews/ci_23198163/colorado-senate-gives-final-approval-historic-marijuana-bills.
- Jacobson, M., Chang, T., Anderson, J. M., MacDonald, J., Bluthenthal, R. N., & Ashwood, S. C. (2011). Regulating medical marijuana dispensaries: An overview with preliminary evidence of their impact on crime. RAND Technical Report, TR-987-RWJ. Santa Monica, CA: RAND Corporation.
- Johnson, G. (2013). Recreational marijuana bars test limits of legal weed laws in Washington, Colorado. *Huffington Post*. Retrieved September 24, 2013, from http://www.huffingtonpost.com/2013/03/26/recreational-marijuana-ba_n_2955049.html.

- Jones, J. M. (2008). Beer back to double-digit lead over wine as favored drink. Gallup. Retrieved September 24, 2013, from <http://www.gallup.com/poll/109066/beer-back-doubledigit-lead-over-wine-favored-drink.aspx>.
- Kelly, E., & Rasul, I. (2012). Policing cannabis and drug related hospital admissions: Evidence from administrative records. Working paper, Department of Economics, University College London. Retrieved November 7, 2012, from <http://www.ucl.ac.uk/~uctpimr/research/>.
- Kelly, E., Darke, S., & Ross, J. (2004). A review of drug use and driving: Epidemiology, impairment, risk factors and risk perceptions. *Drug and Alcohol Review*, 23, 319–344.
- Livingston, B. (2013). Don't call him the "Pot Czar". *The Stranger*, April 10. Retrieved September 24, 2013, from <http://www.thestranger.com/seattle/dont-call-him-the-pot-czar/Content?oid=16460452>.
- Lynne-Landsman, S. D., Livingston, M. D., & Wagenaar, A. C. (2013). Effects of state medical marijuana laws on adolescent marijuana use. *American Journal of Public Health*, 103, 1500–1506.
- Marczinski, C., Harrison, E., & Fillmore, M. (2008). Effects of alcohol on simulated driving and perceived driving impairment in binge drinkers. *Alcoholism: Clinical and Experimental Research*, 32, 1329–1337.
- Markowitz, S. (2000). The price of alcohol, wife abuse, and husband abuse. *Southern Economic Journal*, 67, 279–303.
- Markowitz, S. (2005). Alcohol, drugs and violent crime. *International Review of Law and Economics*, 25, 20–44.
- Markowitz, S., & Grossman, M. (1998). Alcohol regulation and domestic violence towards children. *Contemporary Economic Policy*, 16, 309–320.
- Melberg, H. O., Jones, A. M., & Bretteville-Jensen, A. L. (2010). Is cannabis a gateway to hard drugs? *Empirical Economics*, 38, 583–560.
- Meyer, J., & Glover, S. (2009). Medical marijuana dispensaries will no longer be prosecuted, U.S. Attorney General says. *Los Angeles Times*. Retrieved September 24, 2013, from <http://articles.latimes.com/2009/mar/19/local/me-medpot19>.
- Miron, J. A., & Zwiebel, J. (1995). The economic case against drug prohibition. *Journal of Economic Perspectives*, 9, 175–192.
- Model, K. E. (1993). The effect of marijuana decriminalization on hospital emergency room drug episodes: 1975–1978. *Journal of the American Statistical Association*, 88, 737–747.
- Montgomery, M. (2010). Plummeting marijuana prices create a panic in California. *National Public Radio*, May 15. Retrieved September 24, 2013, from <http://www.npr.org/templates/story/story.php?storyId=126806429>.
- Nagourney, A. (2012). Marijuana, not yet legal for Californians, might as well be. *New York Times*. Retrieved September 24, 2013, from http://www.nytimes.com/2012/12/21/us/politics/stigma-fading-marijuana-common-in-california.html?_r=0.
- National Survey on Drug Use and Health. (2012). State estimates of substance use. Retrieved September 24, 2013, from <https://nsduhweb.rti.org/>.
- Nisbet, C., & Vakil, F. (1972). Some estimates of the price and expenditure elasticities of demand for marijuana among U.C.L.A. students. *Review of Economics and Statistics*, 54, 473–475.
- Pacula, R. L. (1998). Does increasing the beer tax reduce marijuana consumption? *Journal of Health Economics*, 17, 557–586.
- Pacula, R. L., Grossman, M., Chaloupka, F. J., O'Malley, P. M., Johnston, L. D., & Farrelly, M. C. (2001). Marijuana and youth. In J. Gruber (Ed.), *Risky behavior among youths: An economic analysis* (pp. 271–326). Chicago, IL: University of Chicago Press.
- Pacula, R. L., Kilmer, B., Grossman, M., & Chaloupka, F. (2010). Risks and prices: The role of user sanctions in marijuana markets. *B.E. Journal of Economic Analysis and Policy*, 10, 1–36.
- Peck, R. C., Gebers, M. A., Voas, R. B., & Romano, E. (2008). The relationship between blood alcohol concentration (BAC), age, and crash risk. *Journal of Safety Research*, 39, 311–319.

- Robbe, H., & O'Hanlon, J. (1993). Marijuana and actual driving performance. Washington, DC: U.S. Department of Transportation, National Highway Traffic Safety Administration.
- Ronen, A., Gershon, P., Drobiner, H., Rabinovich, A., Bar-Hamburger, R., Mechoulam, R., Yair Cassuto, Y., & Shinar, D. (2008). Effects of THC on driving performance, physiological state and subjective feelings relative to alcohol. *Accident Analysis and Prevention*, 40, 926–934.
- Sewell, R. A., Poling, J., & Sofuoglu, M. (2009). The effect of cannabis compared with alcohol on driving. *American Journal on Addictions*, 18, 185–193.
- Slevin, C. (2010). Colorado medical marijuana raids need to stop: State senators. *Huffington Post*. Retrieved September 24, 2013, from http://www.huffingtonpost.com/2010/03/08/colorado-medical-marijuan_18_n_490519.html.
- Thurstone, C., Lieberman, S. A., & Schmiede, S. J. (2011). Medical marijuana diversion and associated problems in adolescent substance treatment. *Drug Alcohol Dependence*, 118, 489–492.
- Salomonsen-Sautel, S., Sakai, J. T., Thurstone, C., Corley, R., & Hopfer, C. (2012). Medical marijuana use among adolescents in substance abuse treatment. *Journal of the American Academy of Child and Adolescent Psychiatry*, 51, 694–702.
- van Ours, J. C. (2003). Is cannabis a stepping-stone for cocaine? *Journal of Health Economics*, 22, 539–554.
- Williams, J., Pacula, R. L., Chaloupka, F. J., & Wechsler, H. (2004). Alcohol and marijuana use among college students: Economic complements or substitutes? *Health Economics*, 13, 825–843.
- Yörük, B., & Yörük, C. (2011). The impact of minimum legal drinking age laws on alcohol consumption and marijuana use: Evidence from a regression discontinuity design using exact date of birth. *Journal of Health Economics*, 30, 740–753.